

Cite as: A. B. Rosenberg *et al.*, *Science*
10.1126/science.aam8999 (2018).

Single-cell profiling of the developing mouse brain and spinal cord with split-pool barcoding

Alexander B. Rosenberg,^{1,*†} Charles M. Roco,^{2*} Richard A. Muscat,¹ Anna Kuchina,¹ Paul Sample,¹ Zizhen Yao,³ Lucas Gray,³ David J. Peeler,² Sumit Mukherjee,¹ Wei Chen,⁴ Suzie H. Pun,² Drew L. Sellers,^{2,5} Bosiljka Tasic,³ Georg Seelig^{1,4,6†}

¹Department of Electrical Engineering, University of Washington, Seattle, WA, USA. ²Department of Bioengineering, University of Washington, Seattle, WA, USA. ³Allen Institute for Brain Science, Seattle, WA, USA. ⁴Molecular Engineering and Sciences Institute, University of Washington, Seattle, WA, USA. ⁵Institute for Stem Cell and Regenerative Medicine, Seattle, WA, USA. ⁶Paul G. Allen School of Computer Science and Engineering, University of Washington, Seattle, WA, USA.

*These authors contributed equally to this work. †Corresponding author. Email: alex.b.rosenberg@gmail.com (A.B.R.); gseelig@uw.edu (G.S.)

To facilitate scalable profiling of single cells, we developed split-pool ligation-based transcriptome sequencing (SPLiT-seq), a single-cell RNA-seq (scRNA-seq) method that labels the cellular origin of RNA through combinatorial barcoding. SPLiT-seq is compatible with fixed cells or nuclei, allows efficient sample multiplexing and requires no customized equipment. We used SPLiT-seq to analyze 156,049 single-nucleus transcriptomes from postnatal day 2 and 11 mouse brains and spinal cords. Over 100 cell types were identified, with gene expression patterns corresponding to cellular function, regional specificity, and stage of differentiation. Pseudotime analysis revealed transcriptional programs driving four developmental lineages, providing a snapshot of early postnatal development in the murine central nervous system. SPLiT-seq provides a path toward comprehensive single-cell transcriptomic analysis of other similarly complex multicellular systems.

Over 300 years have passed since Leeuwenhoek first described living cells, yet we still do not have a complete catalog of cell types or their functions. Recently, transcriptomic profiling of individual cells has emerged as an essential tool for characterizing cellular diversity (1–3). Single-cell RNA-seq (scRNA-seq) methods have profiled tens of thousands of individual cells (4–6), revealing new insights about cell types within both healthy (7–14) and diseased tissues (15–18). Unfortunately, since these methods require cell sorters, custom microfluidics, or microwells, throughput is still limited and experiments are costly. We introduce SPLiT-seq, a low-cost, scRNA-seq method that enables transcriptional profiling of hundreds of thousands of fixed cells or nuclei in a single experiment. SPLiT-seq does not require partitioning single cells into individual compartments (droplets, microwells or wells), but relies on the cells themselves as compartments. The entire workflow before sequencing consists just of pipetting steps and no complex instruments are needed.

In SPLiT-seq, individual transcriptomes are uniquely labeled by passing a suspension of formaldehyde fixed cells or nuclei through four rounds of combinatorial barcoding. In the first round of barcoding, cells are distributed into a 96-well plate and cDNA is generated with an in-cell reverse transcription (RT) reaction using well-specific barcoded primers. Each well can contain a different biological sample – thereby enabling multiplexing of up to 96 samples in a single experiment. After this step, cells from all wells are

pooled and redistributed into a new 96-well plate, where an in-cell ligation reaction appends a second well-specific barcode to the cDNA. The third-round barcode, which also contains a unique molecular identifier (UMI), is then appended with another round of pooling, splitting, and ligation. After three rounds of barcoding, the cells are pooled, split into sublibraries, and sequencing barcodes are introduced by PCR. This final step provides a fourth barcode, while also making it possible to sequence different numbers of cells in each sublibrary. After sequencing, each transcriptome is assembled by combining reads containing the same four-barcode combination (Fig. 1A and fig. S1A).

Four rounds of combinatorial barcoding can yield 21,233,664 barcode combinations (three rounds of barcoding in 96-well plates followed by a fourth round with 24 PCR reactions) – enough to uniquely label over 1 million cells. Even larger numbers of barcode combinations can be achieved by performing experiments in 384-well plates or through additional rounds of barcoding (fig. S1B). In addition, by performing the first step in a 384-well plate, up to 384 different biological samples could be combined in a single experiment.

SPLiT-seq validation

To test SPLiT-seq's ability to generate uniquely barcoded cells (UBCs), we performed a species-mixing experiment. We mixed cells from one mouse and two human cell lines (NIH/3T3, HEK293, and Hela-S3), fixed them, and used

SPLiT-seq to generate a scRNA-seq library with 1,758 UBCs. The library was sequenced and reads were aligned to a combined mouse-human genome. 99.9% of the UBCs were unambiguously assigned to a single species (>90% of reads aligned to a single genome) with the remaining 0.1% of UBCs representing barcode collisions between mouse and human cells (Fig. 1B). At saturating read coverage (>500,000 reads per cell), we identified a median of 15,365 UMIs and 5,498 genes per human cell and 12,243 UMIs and 4,497 genes per mouse cell. The species purity in both human and mouse UBCs was high: 99.6% of reads in human UBCs and 99.0% of reads in mouse UBCs aligned to their respective genomes. We repeated SPLiT-seq experiments with freshly prepared nuclei as well as nuclei and cells that had been preserved at -80°C for two weeks. In all samples, we detected similar numbers of transcripts and genes per cell (Fig. 1C, fig. S2, and table S1). Gene expression was highly correlated between preserved and freshly prepared cells (Fig. 1D and fig. S2, Pearson-r: 0.987) as well as between cells and nuclei (fig. S2, Pearson-r: 0.952). We also examined gene and UMI detection at different sequencing depths and found that the sensitivity of SPLiT-seq is comparable to droplet-based scRNA-seq methods (fig. S3).

snRNA-seq of developing mouse brain and spinal cord

We used SPLiT-seq to profile nuclei from the developing brain and spinal cord of postnatal day 2 and 11 (P2 and P11) mice. The first round of barcoding assigned identifiers for the P2 brain, P2 spinal cord, P11 brain, and P11 spinal cord samples (Fig. 2A and fig. S4). In total, four rounds of barcoding ($48 \times 96 \times 96 \times 14$) generated over 6 million distinct barcode combinations, making it possible to process hundreds of thousands of nuclei in a single experiment with minimal barcode collisions (2.5% expected collisions for 150,000 nuclei).

To test how many transcripts SPLiT-seq detects within nuclei from the central nervous system, we performed deep sequencing on a sublibrary containing only 131 nuclei. We detected 4,943 UMIs and 2,055 genes per nucleus (UMI duplication: 95%). We then sequenced the rest of the library at lower depth, resulting in a median of 677 genes and 1,022 UMIs per nucleus (UMI duplication: 58%) (table S2). Low-quality transcriptomes were removed from analysis (19), yielding 156,049 single-nucleus transcriptomes (74,862 P2 brain; 7,028 P2 spinal cord; 58,573 P11 brain; 15,586 P11 spinal cord).

Unsupervised clustering grouped transcriptomes into 73 distinct clusters (19) (tables S3 to S5), which were visualized by t-distributed stochastic neighbor embedding (t-SNE) (Fig. 2A). Each of these 73 clusters was assigned to a cell class on the basis of expression of established marker genes (Fig. 2B). Neurons accounted for 79% of the profiled transcrip-

tomies (54 clusters), with most clusters expressing *Meg3*.

The 27,096 non-neuronal transcriptomes spanned 19 different clusters, each assigned to a specific cell type. Four astrocyte types (Fig. 2C) accounted for 50% of all non-neuronal nuclei ($n = 13,481$). Oligodendrocytes (6 types, $n = 4,294$) and oligodendrocyte precursor cells (OPC, 1 type, $n = 5,793$) formed the second most abundant population. We further identified two vascular and leptomeningeal cell (VLMC) types (fig. S5A), endothelial cells, smooth muscle cells (fig. S5B), microglia, macrophages (fig. S5C) (20, 21), ependymal cells, and olfactory ensheathing cells.

Previous work has observed that t-SNE can order cells in 2D space according to stages of differentiation (9). Moving through t-SNE space along the path of differentiation can then be viewed as moving through “pseudotime” (22). As oligogenesis spans the first two postnatal weeks of murine development (23), we asked whether the oligodendrocyte and OPC clusters might reflect a continuous developmental trajectory. When we examined the oligodendrocyte clusters, we found that they formed an overlapping elongated shape in the t-SNE visualization. OPCs and oligodendrocytes from the P2 mouse, were enriched at one end of the structure while oligodendrocytes from the P11 mouse were enriched at the opposite end (fig. S6), indicative of a lineage (19, 22).

We then performed a more thorough analysis of this putative lineage. To ensure that our ordering of oligodendrocytes was determined exclusively by their relationship to other oligodendrocytes, rather than all cells, we re-embedded only transcriptomes within these seven clusters with t-SNE (Fig. 2D and fig. S7A). We calculated the moving average of gene expression in the resulting pseudotime ordering (Fig. 2E and fig. S8). Analysis of these expression patterns confirmed that proliferating OPCs segregated to one end of the t-SNE, whereas mature oligodendrocytes segregated to the opposite end (fig. S7B). We also detected previously reported intermediate stages of oligodendrocyte development, with the order of gene expression across pseudotime nearly identical to the one defined previously (9) (fig. S7C, Spearman-r: 0.94). When analyzing spinal cord and brain derived cells separately, we found more mature oligodendrocytes in the spinal cord than in the brain (fig. S7D), indicating that oligodendrocyte maturation occurs earlier in the spinal cord.

Neuronal cell types

Using known gene markers, we were able to assign most neuronal clusters to specific cell types (19). While some clusters corresponded to abundant cell types, such as cerebellar granule cells (CGCs), others mapped to rare and often less characterized cell types, such as mitral/tufted cells. Previously characterized regional markers were used to assign the majority of clusters to a specific region of the brain (24)

(Fig. 3A). Regional assignments were validated with RNA in situ hybridization (ISH) from the Allen Institute's Developing Mouse Brain Atlas (Allen DMBA) (25). Specifically, we generated composite ISH maps by averaging across the five most highly enriched genes from each of our clusters (tables S6 and S7). For clusters primarily containing P2 or P11 nuclei, we used the P4 or P14 atlases, respectively. The resulting composite maps confirmed the high regional specificity of most types (Fig. 3B and figs. S9 and S10). Cortical pyramidal neuronal types could be further assigned to specific layers using marker genes (Fig. 3C) (7, 8).

Granule cell fate in the hippocampus

In the hippocampus, immature granule cells in the dentate gyrus give rise not only to mature granule cells, but also to pyramidal neurons (26). This process is one of two instances of neurogenesis that continues into adulthood (27), but little is known about the underlying transcriptional program. We determined that three neuronal cell types from the hippocampus likely constituted a developmental trajectory (19). Analysis of only these transcriptomes with t-SNE revealed a clear branching structure (Fig. 3D and fig. S11A). The transcription factor *Prox1*, suspected to be necessary for granule cell identity (28), was exclusively expressed in one branch, whereas genes known to be specific to CA3 pyramidal neurons such as *Spock1* (29) were expressed exclusively in the other branch. Markers of dividing neuronal progenitors were expressed before the branching point and genes in the Slit-Robo signaling pathway were differentially expressed between the two lineages (fig. S11B). We used these data to identify specific temporal dynamics of transcription factors across the two lineages, with *Meis2* as a candidate marker of early pyramidal cell differentiation (Fig. 3E and fig. S12).

Profiling cells in the developing cerebellum

The cerebellum accounts for only 9% of the brain mass in adult mice, but contains nearly 85% of all neurons (30). Despite the wide range of functions performed by the cerebellum, many of the gene expression programs driving development of cerebellar cell types remain unknown. We identified the four main cerebellar neuronal types (Fig. 4A): Purkinje cells, Golgi cells, stellate/basket cells, and CGCs. Two types of Purkinje cells (Fig. 4B) were segregated primarily by age (P2 vs P11), and did not form a continuous trajectory in t-SNE but rather two clearly segregated clusters. The absence of cells at intermediate stages of maturation suggests that Purkinje cell development may be more synchronous than other processes of neurogenesis captured by our dataset.

CGCs, the most numerous type of neuron in the brain (31), drive the postnatal foliation of the cerebellar cortex by migrating from the external granule layer (EGL), through

the molecular layer (ML) and the Purkinje cell layer (PcL) to the internal granule layer (IGL) (32, 33). We created a pseudotime ordering of 15,360 CGCs (Fig. 4C and fig. S13) and measured gene expression across this lineage. We defined genes with specific expression at different points in pseudotime (fig. S14), and then used RNA ISH to map these genes to layers of the developing cerebellar cortex. Genes ordered from early-to-late in pseudotime were progressively expressed from outer-to-inner layers, consistent with the known direction of CGC migration (Fig. 4D). Our analysis revealed previously unknown pseudotime and layer specific gene expression patterns within pathways related to axonal development and neuronal migration (fig. S15).

Origins of cerebellar inhibitory interneurons

The question of whether all cerebellar inhibitory interneurons arise from the same progenitor population has been a point of contention (34). Early hypotheses proposed that stellate/basket cells originated from precursors in the EGL, whereas Golgi cell precursors resided in the ventricular epithelium (35). Later evidence indicated that these two interneurons shared a common precursor in the cerebellar white matter (36, 37). However, the molecular profile of the inhibitory neuron lineage in the cerebellum remains largely unknown.

We found a cerebellar inhibitory interneuron lineage (1517 cells) (Fig. 4E and fig. S16A) with a shared progenitor branching into either Golgi or stellate/basket cells (fig. S17). This lineage includes a known precursor cell type expressing *Pax2* (36), but also a previously unknown, earlier precursor expressing *Pax3* (Fig. 4F). RNA ISH analysis suggests that this *Pax3*+ precursor is located deep within the cerebellar white matter. Moreover, we found that stellate/basket cells expressed genes specific to the molecular layer, whereas Golgi cells expressed genes specific to the granule cell layer (Fig. 4F and fig. S18). The distribution of P2 and P11 nuclei within the lineage clearly demonstrated that the maturation of Golgi cells was well underway by P2 and complete by P11 (fig. S16B). In contrast, stellate/basket cells had not begun to differentiate at P2 and were still not fully mature by P11. These results indicate that the same molecularly defined precursor gives rise to two distinct interneurons at different stages of development.

Cell types in the developing spinal cord

The original clustering was dominated by cells in the brain, and many spinal cord cells did not segregate into well-defined clusters (fig. S19). To resolve more cell types in the spinal cord, we selected all the nuclei originating from the spinal cord and re-clustered them (19), resulting in 44 clusters: 14 non-neuronal types (12 of which were also found in the brain) and 30 neuronal types (Fig. 5A and tables S8 to

S10). We identified 11 different types of GABAergic neurons, of which several were also glycinergic (Fig. 5B). One GABAergic type was identified as cerebrospinal fluid-contacting neurons (CSF-cNs) (38), with the other ten types corresponding to inhibitory interneurons. Glutamatergic interneurons accounted for 15 additional types. We also identified two clusters of cholinergic motor neuron types (alpha and gamma) (39). To date, known markers exist only for gamma motor neurons (*Esrrg* and *Esrrb*) (40), however, we identified specific markers for both alpha and gamma neurons (Fig. 5C).

To infer the spatial origin of neuronal types in the spinal cord, we identified the ten most enriched genes in each type according to our snRNA-seq data and created composite ISH maps based on the Allen Mouse Spinal Cord Atlas (41) (Fig. 5D and fig. S20). Some interneuron subtypes appeared to originate primarily from laminae 1-3, with others originating from laminae 4-6. We found both inhibitory and excitatory neurons in each region. Motor neurons expressed genes found in laminae 9, while CSF-cNs were the only neuronal type expressing genes found in the central canal. These data allowed us to create an atlas of gene expression in the early spinal cord, providing a rich resource for further understanding development of the central nervous system.

Discussion

In this work, we profiled hundreds of thousands of cells using only basic laboratory equipment with a library preparation cost of ~\$0.01 per cell (fig. S21 and table S11). In our analysis of more than 150,000 single-nucleus transcriptomes from two early postnatal stages, we identified 69 types of cells in the brain and 44 types in the spinal cord. We defined many new molecular markers for specific cell types, and explored gene expression in four different developmental lineages.

SPLiT-seq's compatibility with fixed cells and fixed nuclei overcomes challenges faced by other scRNA-seq methods. Fixation can reduce perturbations to endogenous gene expression during cell handling (42) and makes it possible to store cells for future experiments. Moreover, the use of nuclei bypasses the need to obtain intact single cells, which can be challenging for many complex tissues. SPLiT-seq's compatibility with formaldehyde-fixed nuclei suggests it may be used to profile single nuclei from formalin-fixed, paraffin-embedded tissue (43).

SPLiT-seq enables flexible and scalable cell and sample multiplexing. The use of the first-round barcode as a sample identifier makes it possible to profile a large number and variety of samples in parallel, thus minimizing batch effects. As the number of unique barcodes grows exponentially with the number of barcoding rounds, larger numbers of cells than presented here could be processed by adding a fifth

barcoding round or by switching to a 384-well plate format. Although for such large cell numbers sequencing cost may currently be forbidding, it is easy to imagine extended applications, such as targeted sequencing of gene panels, which would even now benefit from very large cell numbers and only require shallow sequencing depth.

Our hope is that the increased scale and accessibility provided by the low cost and minimal equipment requirements of SPLiT-seq will further accelerate the widespread adoption of scRNA-seq.

REFERENCES AND NOTES

1. S. Picelli, Å. K. Björklund, O. R. Faridani, S. Sagasser, G. Winberg, R. Sandberg, Smart-seq2 for sensitive full-length transcriptome profiling in single cells. *Nat. Methods* **10**, 1096–1098 (2013). [doi:10.1038/nmeth.2639](https://doi.org/10.1038/nmeth.2639) Medline
2. T. Hashimshony, F. Wagner, N. Sher, I. Yanai, CEL-Seq: Single-cell RNA-seq by multiplexed linear amplification. *Cell Reports* **2**, 666–673 (2012). [doi:10.1016/j.celrep.2012.08.003](https://doi.org/10.1016/j.celrep.2012.08.003) Medline
3. D. A. Jaitin, E. Kenigsberg, H. Keren-Shaul, N. Elefant, F. Paul, I. Zaretsky, A. Mildner, N. Cohen, S. Jung, A. Tanay, I. Amit, Massively parallel single-cell RNA-seq for marker-free decomposition of tissues into cell types. *Science* **343**, 776–779 (2014). [doi:10.1126/science.1247651](https://doi.org/10.1126/science.1247651) Medline
4. E. Z. Macosko, A. Basu, R. Satija, J. Nemesh, K. Shekhar, M. Goldman, I. Tirosh, A. R. Bialas, N. Kamitaki, E. M. Martersteck, J. J. Trombetta, D. A. Weitz, J. R. Sanes, A. K. Shalek, A. Regev, S. A. McCarroll, Highly parallel genome-wide expression profiling of individual cells using nanoliter droplets. *Cell* **161**, 1202–1214 (2015). [doi:10.1016/j.cell.2015.05.002](https://doi.org/10.1016/j.cell.2015.05.002) Medline
5. A. M. Klein, L. Mazutis, I. Akartuna, N. Tallapragada, A. Veres, V. Li, L. Peshkin, D. A. Weitz, M. W. Kirschner, Droplet barcoding for single-cell transcriptomics applied to embryonic stem cells. *Cell* **161**, 1187–1201 (2015). [doi:10.1016/j.cell.2015.04.044](https://doi.org/10.1016/j.cell.2015.04.044) Medline
6. G. X. Y. Zheng, J. M. Terry, P. Belgrader, P. Ryvkin, Z. W. Bent, R. Wilson, S. B. Ziraldo, T. D. Wheeler, G. P. McDermott, J. Zhu, M. T. Gregory, J. Shuga, L. Montesclaros, J. G. Underwood, D. A. Masquelier, S. Y. Nishimura, M. Schnall-Levin, P. W. Wyatt, C. M. Hindson, R. Bharadwaj, A. Wong, K. D. Ness, L. W. Beppu, H. J. Deeg, C. McFarland, K. R. Loeb, W. J. Valente, N. G. Ericson, E. A. Stevens, J. P. Radich, T. S. Mikkelsen, B. J. Hindson, J. H. Bielas, Massively parallel digital transcriptional profiling of single cells. *Nat. Commun.* **8**, 14049 (2017). [doi:10.1038/ncomms14049](https://doi.org/10.1038/ncomms14049) Medline
7. B. Tasic, V. Menon, T. N. Nguyen, T. K. Kim, T. Jarsky, Z. Yao, B. Levi, L. T. Gray, S. A. Sorensen, T. Dolbeare, D. Bertagnolli, J. Goldy, N. Shapovalova, S. Parry, C. Lee, K. Smith, A. Bernard, L. Madisen, S. M. Sunkin, M. Hawrylycz, C. Koch, H. Zeng, Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. *Nat. Neurosci.* **19**, 335–346 (2016). [doi:10.1038/nn.4216](https://doi.org/10.1038/nn.4216) Medline
8. A. Zeisel, A. B. Muñoz-Manchado, S. Codeluppi, P. Lönnerberg, G. La Manno, A. Jureus, S. Marques, H. Munguba, L. He, C. Betsholtz, C. Rolny, G. Castelo-Branco, J. Hjerling-Leffler, S. Linnarsson, Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq. *Science* **347**, 1138–1142 (2015). [doi:10.1126/science.aaa1934](https://doi.org/10.1126/science.aaa1934) Medline
9. S. Marques, A. Zeisel, S. Codeluppi, D. van Bruggen, A. Mendanha Falcão, L. Xiao, H. Li, M. Häring, H. Hochgerner, R. A. Romanov, D. Gyllborg, A. Muñoz-Manchado, G. La Manno, P. Lönnerberg, E. M. Floridia, F. Rezayee, P. Ernfors, E. Arenas, J. Hjerling-Leffler, T. Harkany, W. D. Richardson, S. Linnarsson, G. Castelo-Branco, Oligodendrocyte heterogeneity in the mouse juvenile and adult central nervous system. *Science* **352**, 1326–1329 (2016). [doi:10.1126/science.aaf6463](https://doi.org/10.1126/science.aaf6463) Medline
10. S. Darmanis, S. A. Sloan, Y. Zhang, M. Enge, C. Caneda, L. M. Shuer, M. G. Hayden Gephart, B. A. Barres, S. R. Quake, A survey of human brain transcriptome diversity at the single cell level. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 7285–7290 (2015). [doi:10.1073/pnas.1507125112](https://doi.org/10.1073/pnas.1507125112) Medline
11. B. B. Lake, R. Ai, G. E. Kaeser, N. S. Salathia, Y. C. Yung, R. Liu, A. Wildberg, D. Gao, H.-L. Fung, S. Chen, R. Vijayaraghavan, J. Wong, A. Chen, X. Sheng, F. Kaper, R. Shen, M. Ronaghi, J.-B. Fan, W. Wang, J. Chun, K. Zhang, Neuronal

- subtypes and diversity revealed by single-nucleus RNA sequencing of the human brain. *Science* **352**, 1586–1590 (2016). [doi:10.1126/science.aaf1204](https://doi.org/10.1126/science.aaf1204) [Medline](#)
12. A. K. Shalek, R. Satija, X. Adiconis, R. S. Gertner, J. T. Gaubloome, R. Raychowdhury, S. Schwartz, N. Yosef, C. Malboeuf, D. Lu, J. J. Trombetta, D. Gennert, A. Gnirke, A. Goren, N. Hacohen, J. Z. Levin, H. Park, A. Regev, Single-cell transcriptomics reveals bimodality in expression and splicing in immune cells. *Nature* **498**, 236–240 (2013). [doi:10.1038/nature12172](https://doi.org/10.1038/nature12172) [Medline](#)
 13. D. Grün, A. Lyubimova, L. Kester, K. Wiebrands, O. Basak, N. Sasaki, H. Clevers, A. van Oudenaarden, Single-cell messenger RNA sequencing reveals rare intestinal cell types. *Nature* **525**, 251–255 (2015). [doi:10.1038/nature14966](https://doi.org/10.1038/nature14966) [Medline](#)
 14. V. Moignard, S. Woodhouse, L. Haghverdi, A. J. Lilly, Y. Tanaka, A. C. Wilkinson, F. Büttner, I. C. Macaulay, W. Jawaid, E. Diamanti, S.-I. Nishikawa, N. Piterman, V. Kouskoff, F. J. Theis, J. Fisher, B. Göttgens, Decoding the regulatory network of early blood development from single-cell gene expression measurements. *Nat. Biotechnol.* **33**, 269–276 (2015). [doi:10.1038/nbt.3154](https://doi.org/10.1038/nbt.3154) [Medline](#)
 15. A. S. Venteicher, I. Tirosh, C. Hebert, K. Yizhak, C. Neftel, M. G. Filbin, V. Hovestadt, L. E. Escalante, M. L. Shaw, C. Rodman, S. M. Gillespie, D. Dionne, C. C. Luo, H. Ravichandran, R. Mylvaganam, C. Mount, M. L. Onozato, B. V. Nahed, H. Wakimoto, W. T. Curry, A. J. Iafrate, M. N. Rivera, M. P. Frosch, T. R. Golub, P. K. Brastianos, G. Getz, A. P. Patel, M. Monje, D. P. Cahill, O. Rozenblatt-Rosen, D. N. Louis, B. E. Bernstein, A. Regev, M. L. Suvà, Decoupling genetics, lineages, and microenvironment in IDH-mutant gliomas by single-cell RNA-seq. *Science* **355**, eaai8478 (2017). [doi:10.1126/science.aai8478](https://doi.org/10.1126/science.aai8478) [Medline](#)
 16. I. Tirosh, B. Izar, S. M. Prakadan, M. H. Wadsworth 2nd, D. Treacy, J. J. Trombetta, A. Rotem, C. Rodman, C. Lian, G. Murphy, M. Fallahi-Sichani, K. Dutton-Regester, J.-R. Lin, O. Cohen, P. Shah, D. Lu, A. S. Genshaft, T. K. Hughes, C. G. K. Ziegler, S. W. Kazer, A. Gaillard, K. E. Kolb, A.-C. Villani, C. M. Johannessen, A. Y. Andreev, E. M. Van Allen, M. Bertagnolli, P. K. Sorger, R. J. Sullivan, K. T. Flaherty, D. T. Frederick, J. Jané-Valbuena, C. H. Yoon, O. Rozenblatt-Rosen, A. K. Shalek, A. Regev, L. A. Garraway, Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science* **352**, 189–196 (2016). [doi:10.1126/science.aad0501](https://doi.org/10.1126/science.aad0501) [Medline](#)
 17. L. Sang, H. A. Collier, J. M. Roberts, Control of the reversibility of cellular quiescence by the transcriptional repressor HES1. *Science* **321**, 1095–1100 (2008). [doi:10.1126/science.1155998](https://doi.org/10.1126/science.1155998) [Medline](#)
 18. C. Zheng, L. Zheng, J.-K. Yoo, H. Guo, Y. Zhang, X. Guo, B. Kang, R. Hu, J. Y. Huang, Q. Zhang, Z. Liu, M. Dong, X. Hu, W. Ouyang, J. Peng, Z. Zhang, Landscape of infiltrating T cells in liver cancer revealed by single-cell sequencing. *Cell* **169**, 1342–1356.e16 (2017). [doi:10.1016/j.cell.2017.05.035](https://doi.org/10.1016/j.cell.2017.05.035) [Medline](#)
 19. Materials and methods are provided as supplementary materials.
 20. S. E. Hickman, N. D. Kingery, T. K. Ohsumi, M. L. Borowsky, L. C. Wang, T. K. Means, J. El Khoury, The microglial sensome revealed by direct RNA sequencing. *Nat. Neurosci.* **16**, 1896–1905 (2013). [doi:10.1038/nn.3554](https://doi.org/10.1038/nn.3554) [Medline](#)
 21. O. Matcovitch-Natan, D. R. Winter, A. Giladi, S. Vargas Aguilar, A. Spinrad, S. Sarrazin, H. Ben-Yehuda, E. David, F. Zelada González, P. Perrin, H. Keren-Shaul, M. Gury, D. Lara-Astaiso, C. A. Thaiss, M. Cohen, K. Bahar Halpern, K. Baruch, A. Deczkowska, E. Lorenzo-Vivas, S. Itzkovitz, E. Elinav, M. H. Sieveke, M. Schwartz, I. Amit, Microglia development follows a stepwise program to regulate brain homeostasis. *Science* **353**, aad8670 (2016). [doi:10.1126/science.aad8670](https://doi.org/10.1126/science.aad8670) [Medline](#)
 22. C. Trapnell, D. Cacchiarelli, J. Grimsby, P. Pokharel, S. Li, M. Morse, N. J. Lennon, K. J. Livak, T. S. Mikkelsen, J. L. Rinn, The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat. Biotechnol.* **32**, 381–386 (2014). [doi:10.1038/nbt.2859](https://doi.org/10.1038/nbt.2859) [Medline](#)
 23. S. W. Levison, J. E. Goldman, Both oligodendrocytes and astrocytes develop from progenitors in the subventricular zone of postnatal rat forebrain. *Neuron* **10**, 201–212 (1993). [doi:10.1016/0896-6273\(93\)90311-F](https://doi.org/10.1016/0896-6273(93)90311-F) [Medline](#)
 24. E. S. Lein, M. J. Hawrylycz, N. Ao, M. Ayres, A. Bensinger, A. Bernard, A. F. Boe, M. S. Boguski, K. S. Brockway, E. J. Byrnes, L. Chen, L. Chen, T.-M. Chen, M. C. Chin, J. Chong, B. E. Crook, A. Czaplinska, C. N. Dang, S. Datta, N. R. Dee, A. L. Desaki, T. Desta, E. Diep, T. A. Dolbeare, M. J. Donelan, H.-W. Dong, J. G. Dougherty, B. J. Duncan, A. J. Ebbert, G. Eichele, L. K. Estlin, C. Faber, B. A. Facer, R. Fields, S. R. Fischer, T. P. Fliss, C. Frensley, S. N. Gates, K. J. Glatfelter, K. R. Halverson, M. R. Hart, J. G. Hohmann, M. P. Howell, D. P. Jeung, R. A. Johnson, P. T. Karr, R. Kaval, J. M. Kidney, R. H. Knapik, C. L. Kuan, J. H. Lake, A. R. Laramée, K. D. Larsen, C. Lau, T. A. Lemon, A. J. Liang, Y. Liu, L. T. Luong, J. Michaels, J. J. Morgan, R. J. Morgan, M. T. Mortrud, N. F. Mosqueda, L. L. Ng, R. Ng, G. J. Orta, C. C. Overly, T. H. Pak, S. E. Parry, S. D. Pathak, O. C. Pearson, R. B. Puchalski, Z. L. Riley, H. R. Rockett, S. A. Rowland, J. J. Royall, M. J. Ruiz, N. R. Sarno, K. Schaffnit, N. V. Shapovalova, T. S. Visay, C. R. Slaughterbeck, S. C. Smith, K. A. Smith, B. I. Smith, A. J. Sotd, N. N. Stewart, K.-R. Stumpf, S. M. Sunkin, M. Sutram, A. Tam, C. D. Teemer, C. Thaller, C. L. Thompson, L. R. Varnam, A. Visel, R. M. Whitlock, P. E. Wornoutka, C. K. Wolkey, V. Y. Wong, M. Wood, M. B. Yaylaoglu, R. C. Young, B. L. Youngstrom, X. F. Yuan, B. Zhang, T. A. Zwingman, A. R. Jones, Genome-wide atlas of gene expression in the adult mouse brain. *Nature* **445**, 168–176 (2007). [doi:10.1038/nature05453](https://doi.org/10.1038/nature05453) [Medline](#)
 25. Allen Institute for Brain Science, Developing Mouse Brain Atlas (2008), (available at <http://developingmouse.brain-map.org/>).
 26. T. Iwano, A. Masuda, H. Kiyonari, H. Enomoto, F. Matsuzaki, Prox1 postmitotically defines dentate gyrus cells by specifying granule cell identity over CA3 pyramidal cell fate in the hippocampus. *Development* **139**, 3051–3062 (2012). [doi:10.1242/dev.080002](https://doi.org/10.1242/dev.080002) [Medline](#)
 27. C. Zhao, W. Deng, F. H. Gage, Mechanisms and functional implications of adult neurogenesis. *Cell* **132**, 645–660 (2008). [doi:10.1016/j.cell.2008.01.033](https://doi.org/10.1016/j.cell.2008.01.033) [Medline](#)
 28. A. Lavado, O. V. Lagutin, L. M. L. Chow, S. J. Baker, G. Oliver, Prox1 is required for granule cell maturation and intermediate progenitor maintenance during brain neurogenesis. *PLOS Biol.* **8**, e1000460 (2010). [doi:10.1371/journal.pbio.1000460](https://doi.org/10.1371/journal.pbio.1000460) [Medline](#)
 29. F. Bonnet, J.-P. Périn, F. Charbonnier, A. Camuzat, G. Roussel, J.-L. Nussbaum, P. M. Alliel, Structure and cellular distribution of mouse brain testican. Association with the postsynaptic area of hippocampus pyramidal cells. *J. Biol. Chem.* **271**, 4373–4380 (1996). [doi:10.1074/jbc.271.8.4373](https://doi.org/10.1074/jbc.271.8.4373) [Medline](#)
 30. S. Herculano-Houzel, The human brain in numbers: A linearly scaled-up primate brain. *Front. Hum. Neurosci.* **3**, 31 (2009). [doi:10.3389/neuro.09.031.2009](https://doi.org/10.3389/neuro.09.031.2009) [Medline](#)
 31. K. Nakashima, H. Umeshima, M. Kengaku, Cerebellar granule cells are predominantly generated by terminal symmetric divisions of granule cell precursors. *Dev. Dyn.* **244**, 748–758 (2015). [doi:10.1002/dvdy.24276](https://doi.org/10.1002/dvdy.24276) [Medline](#)
 32. A. Sudarov, A. L. Joyner, Cerebellum morphogenesis: The foliation pattern is orchestrated by multi-cellular anchoring centers. *Neural Dev.* **2**, 26 (2007). [doi:10.1186/1749-8104-2-26](https://doi.org/10.1186/1749-8104-2-26) [Medline](#)
 33. J. C. Chang, M. Leung, H. N. Gokozan, P. E. Gygli, F. P. Catacutan, C. Zeisler, J. J. Otero, Mitotic events in cerebellar granule progenitor cells that expand cerebellar surface area are critical for normal cerebellar cortical lamination in mice. *J. Neuropathol. Exp. Neurol.* **74**, 261–272 (2015). [doi:10.1097/NFN.0000000000000171](https://doi.org/10.1097/NFN.0000000000000171) [Medline](#)
 34. K. Schilling, J. Oberdick, F. Rossi, S. L. Baader, Besides Purkinje cells and granule neurons: An appraisal of the cell biology of the interneurons of the cerebellar cortex. *Histochem. Cell Biol.* **130**, 601–615 (2008). [doi:10.1007/s00418-008-0483-y](https://doi.org/10.1007/s00418-008-0483-y) [Medline](#)
 35. J. Altman, S. A. Bayer, Development of the precerebellar nuclei in the rat: I. The precerebellar neuroepithelium of the rhombencephalon. *J. Comp. Neurol.* **257**, 477–489 (1987). [doi:10.1002/cne.902570402](https://doi.org/10.1002/cne.902570402) [Medline](#)
 36. S. M. Maricich, K. Herrup, Pax-2 expression defines a subset of GABAergic interneurons and their precursors in the developing murine cerebellum. *J. Neurobiol.* **41**, 281–294 (1999). [doi:10.1002/\(SICI\)1097-4695\(19991105\)41:2<281::AID-NEU10>3.0.CO;2-5](https://doi.org/10.1002/(SICI)1097-4695(19991105)41:2<281::AID-NEU10>3.0.CO;2-5) [Medline](#)
 37. G. Weisheit, M. Gliem, E. Endl, P. L. Pfeffer, M. Busslinger, K. Schilling, Postnatal development of the murine cerebellar cortex: Formation and early dispersal of basket, stellate and Golgi neurons. *Eur. J. Neurosci.* **24**, 466–478 (2006). [doi:10.1111/j.1460-9568.2006.04915.x](https://doi.org/10.1111/j.1460-9568.2006.04915.x) [Medline](#)
 38. Y. L. Petracca, M. M. Sartoretti, D. J. Di Bella, A. Marin-Burgin, A. L. Carcagno, A. F. Schinder, G. M. Lanuza, The late and dual origin of cerebrospinal fluid-contacting neurons in the mouse spinal cord. *Development* **143**, 880–891 (2016). [doi:10.1242/dev.129254](https://doi.org/10.1242/dev.129254) [Medline](#)
 39. A. Enjin, N. Rabe, S. T. Nakanishi, A. Vallstedt, H. Gezelius, F. Memic, M. Lind, T. Hjält, W. G. Tourtellotte, C. Bruder, G. Eichele, P. J. Whelan, K. Kullander, Identification of novel spinal cholinergic genetic subtypes disclose Chodl and Ptx2 as markers for fast motor neurons and partition cells. *J. Comp. Neurol.* **518**,

- 2284–2304 (2010). [doi:10.1002/cne.22332](https://doi.org/10.1002/cne.22332) [Medline](#)
40. M. Lalancette-Hebert, A. Sharma, A. K. Lyashchenko, N. A. Shneider, Gamma motor neurons survive and exacerbate alpha motor neuron degeneration in ALS. *Proc. Natl. Acad. Sci. U.S.A.* **113**, E8316–E8325 (2016). [doi:10.1073/pnas.1605210113](https://doi.org/10.1073/pnas.1605210113) [Medline](#)
 41. Allen Institute for Brain Science, Allen Mouse Spinal Cord Atlas (2008); available at <http://mousespinal.brain-map.org/imageseries/showref.html>.
 42. B. Lacar, S. B. Linker, B. N. Jaeger, S. Krishnaswami, J. Barron, M. Kelder, S. Parylak, A. Paquola, P. Venepally, M. Novotny, C. O'Connor, C. Fitzpatrick, J. Erwin, J. Y. Hsu, D. Husband, M. J. McConnell, R. Lasken, F. H. Gage, Nuclear RNA-seq of single neurons reveals molecular signatures of activation. *Nat. Commun.* **7**, 11022 (2016). [doi:10.1038/ncomms11022](https://doi.org/10.1038/ncomms11022) [Medline](#)
 43. E. R. Thomsen, J. K. Mich, Z. Yao, R. D. Hodge, A. M. Doyle, S. Jang, S. I. Shehata, A. M. Nelson, N. V. Shapovalova, B. P. Levi, S. Ramanathan, Fixed single-cell transcriptomic characterization of human radial glial diversity. *Nat. Methods* **13**, 87–93 (2016). [doi:10.1038/nmeth.3629](https://doi.org/10.1038/nmeth.3629) [Medline](#)
 44. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, T. R. Gingeras, STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013). [doi:10.1093/bioinformatics/bts635](https://doi.org/10.1093/bioinformatics/bts635) [Medline](#)
 45. E. Zorita, P. Cuscó, G. J. Filion, Starcode: Sequence clustering based on all-pairs search. *Bioinformatics* **31**, 1913–1919 (2015). [doi:10.1093/bioinformatics/btv053](https://doi.org/10.1093/bioinformatics/btv053) [Medline](#)
 46. Z. Yao, J. K. Mich, S. Ku, V. Menon, A.-R. Krostag, R. A. Martinez, L. Furchtgott, H. Mulholland, S. Bort, M. A. Fuqua, B. W. Gregor, R. D. Hodge, A. Jayabalu, R. C. May, S. Melton, A. M. Nelson, N. K. Ngo, N. V. Shapovalova, S. I. Shehata, M. W. Smith, L. J. Tait, C. L. Thompson, E. R. Thomsen, C. Ye, I. A. Glass, A. Kaykas, S. Yao, J. W. Phillips, J. S. Grimley, B. P. Levi, Y. Wang, S. Ramanathan, A single-cell roadmap of lineage bifurcation in human ESC models of embryonic brain development. *Cell Stem Cell* **20**, 120–134 (2017). [doi:10.1016/j.stem.2016.09.011](https://doi.org/10.1016/j.stem.2016.09.011) [Medline](#)
 47. K. Shekhar, S. W. Lapan, I. E. Whitney, N. M. Tran, E. Z. Macosko, M. Kowalczyk, X. Adiconis, J. Z. Levin, J. Nemesh, M. Goldman, S. A. McCarroll, C. L. Cepko, A. Regev, J. R. Sanes, Comprehensive classification of retinal bipolar neurons by single-cell transcriptomics. *Cell* **166**, 1308–1323.e30 (2016). [doi:10.1016/j.cell.2016.07.054](https://doi.org/10.1016/j.cell.2016.07.054) [Medline](#)
 48. N. Habib, Y. Li, M. Heidenreich, L. Swiech, I. Avraham-David, J. J. Trombetta, C. Hession, F. Zhang, A. Regev, Div-Seq: Single-nucleus RNA-Seq reveals dynamics of rare adult newborn neurons. *Science* **353**, 925–928 (2016). [doi:10.1126/science.aad7038](https://doi.org/10.1126/science.aad7038) [Medline](#)
 49. F. Pedregosa et al., Scikit-learn: Machine learning in Python. *J. Mach. Learn. Res.* **12**, 2825–2830 (2011).
 50. L. van der Maaten, Accelerating t-SNE using tree-based algorithms. *J. Mach. Learn. Res.* **15**, 3221–3245 (2014).
 51. M. Ester, H.-P. Kriegel, J. Sander, X. Xu, A density-based algorithm for discovering clusters a density-based algorithm for discovering clusters in large spatial databases with noise. *Proc. Second Int. Conf. Knowl. Discov. Data Min.* (1996), pp. 226–231.
 52. X. Qiu, Q. Mao, Y. Tang, L. Wang, R. Chawla, H. A. Pliner, C. Trapnell, Reversed graph embedding resolves complex single-cell trajectories. *Nat. Methods* **14**, 979–982 (2017). [doi:10.1038/nmeth.4402](https://doi.org/10.1038/nmeth.4402) [Medline](#)
 53. M. J. Yang, S. Sim, J. H. Jeon, E. Jeong, H.-C. Kim, Y.-J. Park, I.-B. Kim, Mitral and tufted cells are potential cellular targets of nitration in the olfactory bulb of aged mice. *PLOS ONE* **8**, e59673 (2013). [doi:10.1371/journal.pone.0059673](https://doi.org/10.1371/journal.pone.0059673) [Medline](#)
 54. Y. I. Kawasaki, A. C. Salzberg, M. Li, N. Šestan, C. A. Greer, F. Imamura, RNA-seq analysis of developing olfactory bulb projection neurons. *Mol. Cell. Neurosci.* **74**, 78–86 (2016). [doi:10.1016/j.mcn.2016.03.009](https://doi.org/10.1016/j.mcn.2016.03.009) [Medline](#)
 55. J. P. Doyle, J. D. Dougherty, M. Heiman, E. F. Schmidt, T. R. Stevens, G. Ma, S. Bupp, P. Shrestha, R. D. Shah, M. L. Doughty, S. Gong, P. Greengard, N. Heintz, Application of a translational profiling approach for the comparative analysis of CNS cell types. *Cell* **135**, 749–762 (2008). [doi:10.1016/j.cell.2008.10.029](https://doi.org/10.1016/j.cell.2008.10.029) [Medline](#)
 56. E. A. Alcamo, L. Chirivella, M. Dautzenberg, G. Dobrova, I. Fariñas, R. Grosschedl, S. K. McConnell, Satb2 regulates callosal projection neuron identity in the developing cerebral cortex. *Neuron* **57**, 364–377 (2008). [doi:10.1016/j.neuron.2007.12.012](https://doi.org/10.1016/j.neuron.2007.12.012) [Medline](#)
 57. K. Hesse, K. Vaupel, S. Kurt, R. Buettner, J. Kirfel, M. Moser, AP-2δ is a crucial transcriptional regulator of the posterior midbrain. *PLOS ONE* **6**, e23483 (2011). [doi:10.1371/journal.pone.0023483](https://doi.org/10.1371/journal.pone.0023483) [Medline](#)
 58. H. Valcanis, S.-S. Tan, Layer specification of transplanted interneurons in developing mouse neocortex. *J. Neurosci.* **23**, 5113–5122 (2003). [Medline](#)
 59. M. Ogawa, T. Miyata, K. Nakajima, K. Yagyu, M. Seike, K. Ikenaka, H. Yamamoto, K. Mikoshiba, The reeler gene-associated antigen on Cajal-Retzius neurons is a crucial molecule for laminar organization of cortical neurons. *Neuron* **14**, 899–912 (1995). [doi:10.1016/0896-6273\(95\)90329-1](https://doi.org/10.1016/0896-6273(95)90329-1) [Medline](#)
 60. L. He, M. Vanlandewijck, E. Raschperger, M. Andaloussi Mäe, B. Jung, T. Lebouvier, K. Ando, J. Hofmann, A. Keller, C. Betsholtz, Analysis of the brain mural cell transcriptome. *Sci. Rep.* **6**, 35108 (2016). [doi:10.1038/srep35108](https://doi.org/10.1038/srep35108) [Medline](#)
 61. Y.-C. Chiu, M.-Y. Li, Y.-H. Liu, J.-Y. Ding, J.-Y. Yu, T.-W. Wang, Foxp2 regulates neuronal differentiation and neuronal subtype specification. *Dev. Neurobiol.* **74**, 723–738 (2014). [doi:10.1002/dneu.22166](https://doi.org/10.1002/dneu.22166) [Medline](#)
 62. T. Miyashita, M. Wintzer, T. Kurotani, T. Konishi, N. Ichinohe, K. S. Rockland, Neurotrophin-3 is involved in the formation of apical dendritic bundles in cortical layer 2 of the rat. *Cereb. Cortex* **20**, 229–240 (2010). [doi:10.1093/cercor/bhp093](https://doi.org/10.1093/cercor/bhp093) [Medline](#)
 63. T. G. Belgard, A. C. Marques, P. L. Oliver, H. O. Abaan, T. M. Sirey, A. Hoerder-Suabedissen, F. García-Moreno, Z. Molnár, E. H. Margulies, C. P. Ponting, A transcriptomic atlas of mouse neocortical layers. *Neuron* **71**, 605–616 (2011). [doi:10.1016/j.neuron.2011.06.039](https://doi.org/10.1016/j.neuron.2011.06.039) [Medline](#)
 64. A. Nagalski, L. Puelles, M. Dabrowski, T. Wegierski, J. Kuznicki, M. B. Wisniewska, Molecular anatomy of the thalamic complex and the underlying transcription factors. *Brain Struct. Funct.* **221**, 2493–2510 (2016). [doi:10.1007/s00429-015-1052-5](https://doi.org/10.1007/s00429-015-1052-5) [Medline](#)
 65. M. B. Wisniewska, K. Misztal, W. Michowski, M. Szczot, E. Purta, W. Lesniak, M. E. Klejman, M. Dabrowski, R. K. Filipkowski, A. Nagalski, J. W. Mozrzymas, J. Kuznicki, LEF1/beta-catenin complex regulates transcription of the Cav3.1 calcium channel gene (Cacna1g) in thalamic neurons of the adult brain. *J. Neurosci.* **30**, 4957–4969 (2010). [doi:10.1523/JNEUROSCI.1425-09.2010](https://doi.org/10.1523/JNEUROSCI.1425-09.2010) [Medline](#)
 66. H. Song, B. Lee, D. Pyun, J. Guimera, Y. Son, J. Yoon, K. Baek, W. Wurst, Y. Jeong, Ascl1 and Helt act combinatorially to specify thalamic neuronal identity by repressing Dlx5 activation. *Dev. Biol.* **398**, 280–291 (2015). [doi:10.1016/j.ydbio.2014.12.003](https://doi.org/10.1016/j.ydbio.2014.12.003) [Medline](#)
 67. J. D. Corrales, S. Blaess, E. M. Mahoney, A. L. Joyner, The level of sonic hedgehog signaling regulates the complexity of cerebellar foliation. *Development* **133**, 1811–1821 (2006). [doi:10.1242/dev.02351](https://doi.org/10.1242/dev.02351) [Medline](#)
 68. E. Salero, M. E. Hatten, Differentiation of ES cells into cerebellar neurons. *Proc. Natl. Acad. Sci. U.S.A.* **104**, 2997–3002 (2007). [doi:10.1073/pnas.0610879104](https://doi.org/10.1073/pnas.0610879104) [Medline](#)
 69. C. A. D'Souza, V. Chopra, R. Varhol, Y.-Y. Xie, S. Bohacec, Y. Zhao, L. L. C. Lee, M. Bilenky, E. Portales-Casamar, A. He, W. W. Wasserman, D. Goldowitz, M. A. Marra, R. A. Holt, E. M. Simpson, S. J. M. Jones, Identification of a set of genes showing regionally enriched expression in the mouse brain. *BMC Neurosci.* **9**, 66 (2008). [doi:10.1186/1471-2202-9-66](https://doi.org/10.1186/1471-2202-9-66) [Medline](#)
 70. N. X. Tritsch, J. B. Ding, B. L. Sabatini, Dopaminergic neurons inhibit striatal output through non-canonical release of GABA. *Nature* **490**, 262–266 (2012). [doi:10.1038/nature11466](https://doi.org/10.1038/nature11466) [Medline](#)
 71. M. Moriyama, T. Fukuhara, M. Britschgi, Y. He, R. Narasimhan, S. Villeda, H. Molina, B. T. Huber, M. Holers, T. Wyss-Coray, Complement receptor 2 is expressed in neural progenitor cells and regulates adult hippocampal neurogenesis. *J. Neurosci.* **31**, 3981–3989 (2011). [doi:10.1523/JNEUROSCI.3617-10.2011](https://doi.org/10.1523/JNEUROSCI.3617-10.2011) [Medline](#)
 72. J. V. Nielsen, J. B. Blom, J. Norberg, N. A. Jensen, Zbtb20-induced CA1 pyramidal neuron development and area enlargement in the cerebral midline cortex of mice. *Cereb. Cortex* **20**, 1904–1914 (2010). [doi:10.1093/cercor/bhp261](https://doi.org/10.1093/cercor/bhp261) [Medline](#)
 73. M. S. Cembrowski, L. Wang, K. Sugino, B. C. Shields, N. Spruston, HippoSeq: A comprehensive RNA-seq database of gene expression in hippocampal principal neurons. *eLife* **5**, e14997 (2016). [doi:10.7554/eLife.14997](https://doi.org/10.7554/eLife.14997) [Medline](#)
 74. C. L. Thompson, L. Ng, V. Menon, S. Martinez, C.-K. Lee, K. Glattfelder, S. M.

- Sunkin, A. Henry, C. Lau, C. Dang, R. Garcia-Lopez, A. Martinez-Ferre, A. Pombero, J. L. R. Rubenstein, W. B. Wakeman, J. Hohmann, N. Dee, A. J. Södt, R. Young, K. Smith, T.-N. Nguyen, J. Kidney, L. Kuan, A. Jeromin, A. Kaykas, J. Miller, D. Page, G. Orta, A. Bernard, Z. Riley, S. Smith, P. Wahnoutka, M. J. Hawrylycz, L. Puellas, A. R. Jones, A high-resolution spatiotemporal atlas of gene expression of the developing mouse brain. *Neuron* **83**, 309–323 (2014). doi:10.1016/j.neuron.2014.05.033 Medline
75. Z. Xie, X. Ma, W. Ji, G. Zhou, Y. Lu, Z. Xiang, Y. X. Wang, L. Zhang, Y. Hu, Y.-Q. Ding, W. J. Zhang, Zbtb20 is essential for the specification of CA1 field identity in the developing hippocampus. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 6510–6515 (2010). doi:10.1073/pnas.0912315107 Medline
76. H. M. Knight, R. Walker, R. James, D. J. Porteous, W. J. Muir, D. H. R. Blackwood, B. S. Pickard, GRIK4/KA1 protein expression in human brain and correlation with bipolar disorder risk variant status. *Am. J. Med. Genet. B. Neuropsychiatr. Genet.* **159B**, 21–29 (2012). doi:10.1002/ajmg.b.31248 Medline
77. S. Shah, E. Lubeck, W. Zhou, L. Cai, In situ transcription profiling of single cells reveals spatial organization of cells in the mouse hippocampus. *Neuron* **92**, 342–357 (2016). doi:10.1016/j.neuron.2016.10.001 Medline
78. R. P. Seal, X. Wang, Y. Guan, S. N. Raja, C. J. Woodbury, A. I. Basbaum, R. H. Edwards, Injury-induced mechanical hypersensitivity requires C-low threshold mechanoreceptors. *Nature* **462**, 651–655 (2009). doi:10.1038/nature08505 Medline
79. P. Alifragis, A. Liapi, J. G. Parnavelas, Lhx6 regulates the migration of cortical interneurons from the ventral telencephalon but does not specify their GABA phenotype. *J. Neurosci.* **24**, 5643–5648 (2004). doi:10.1523/JNEUROSCI.1245-04.2004 Medline
80. E. Sánchez-Mendoza, V. Bellver-Landete, J. J. Merino, M. P. González, R. Martínez-Murillo, M. J. Oset-Gasque, Review: Could neurotransmitters influence neurogenesis and neurorepair after stroke? *Neuropathol. Appl. Neurobiol.* **39**, 722–735 (2013). doi:10.1111/nap.12082 Medline
81. H. Abraham, C. G. Pérez-García, G. Meyer, p73 and Reelin in Cajal-Retzius cells of the developing human hippocampal formation. *Cereb. Cortex* **14**, 484–495 (2004). doi:10.1093/cercor/bhh010 Medline
82. K.-K. Cheung, S. C. Mok, P. Rezaie, W. Y. Chan, Dynamic expression of Dab2 in the mouse embryonic central nervous system. *BMC Dev. Biol.* **8**, 76 (2008). doi:10.1186/1471-213X-8-76 Medline
83. O. Butovsky, M. P. Jedrychowski, C. S. Moore, R. Cialic, A. J. Lanser, G. Gabriely, T. Koeglspenger, B. Dake, P. M. Wu, C. E. Doykan, Z. Fanek, L. Liu, Z. Chen, J. D. Rothstein, R. M. Ransohoff, S. P. Gygi, J. P. Antel, H. L. Weiner, Identification of a unique TGF- β -dependent molecular and functional signature in microglia. *Nat. Neurosci.* **17**, 131–143 (2014). doi:10.1038/nn.3599 Medline
84. J. T. Ehmsen, Y. Liu, Y. Wang, N. Paladugu, A. E. Johnson, J. D. Rothstein, S. du Lac, M. P. Mattson, A. Höke, The astrocytic transporter SLC7A10 (Asc-1) mediates glycinergic inhibition of spinal cord motor neurons. *Sci. Rep.* **6**, 35592 (2016). doi:10.1038/srep35592 Medline
85. S. Chuikov, B. P. Levi, M. L. Smith, S. J. Morrison, Prdm16 promotes stem cell maintenance in multiple tissues, partly by regulating oxidative stress. *Nat. Cell Biol.* **12**, 999–1006 (2010). doi:10.1038/ncb2101 Medline
86. Z. H. Afsari, W. M. Renno, E. Abd-El-Basset, Alteration of glial fibrillary acidic proteins immunoreactivity in astrocytes of the spinal cord diabetic rats. *Anat. Rec. Adv. Integr. Anat. Evol. Biol.* **291**, 390–399 (2008). doi:10.1002/ar.20678 Medline
87. A. S. Saab, A. Neumeyer, H. M. Jahn, A. Cupido, A. A. M. Šimek, H.-J. Boele, A. Scheller, K. Le Meur, M. Götz, H. Monyer, R. Sprengel, M. E. Rubio, J. W. Deitmer, C. I. De Zeeuw, F. Kirchhoff, Bergmann glial AMPA receptors are required for fine motor coordination. *Science* **337**, 749–753 (2012). doi:10.1126/science.1221140 Medline
88. A. Honoré, S. Le Corre, C. Derambure, R. Normand, C. Duclos, O. Boyer, J.-P. Marie, N. Guérout, Isolation, characterization, and genetic profiling of subpopulations of olfactory ensheathing cells from the olfactory bulb. *Glia* **60**, 404–413 (2012). doi:10.1002/glia.22274 Medline
89. M. Murthy, S. Bocking, F. Verginelli, S. Stifani, Transcription factor Runx1 inhibits proliferation and promotes developmental maturation in a selected population of inner olfactory nerve layer olfactory ensheathing cells. *Gene* **540**, 191–200 (2014). doi:10.1016/j.gene.2014.02.038 Medline
90. J. D. Cahoy, B. Emery, A. Kaushal, L. C. Foo, J. L. Zamanian, K. S. Christopherson, Y. Xing, J. L. Lubischer, P. A. Krieg, S. A. Krupenko, W. J. Thompson, B. A. Barres, A transcriptome database for astrocytes, neurons, and oligodendrocytes: A new resource for understanding brain development and function. *J. Neurosci.* **28**, 264–278 (2008). doi:10.1523/JNEUROSCI.4178-07.2008 Medline
91. M. G. Erlander, A. J. Tobin, The structural and functional heterogeneity of glutamic acid decarboxylase: A review. *Neurochem. Res.* **16**, 215–226 (1991). doi:10.1007/BF00966084 Medline
92. P. R. Brumovsky, VGLUTs in peripheral neurons and the spinal cord: Time for a review. *ISRN Neurol.* **2013**, 829753 (2013). doi:10.1155/2013/829753 Medline

ACKNOWLEDGMENTS

We would like to thank Thuc Nghi Nguyen for help with cluster identity assignment.

Funding: This work was supported by NIH R01CA207029 and NSF CCF-1317653 to G.S. and NIH R01NS064404 and R21NS086500 to S.H.P., Z.Y., L.G. and B.T. were supported by the Allen Institute for Brain Science. C.M.R. was supported by the National Center for Advancing Translational Sciences of the National Institutes of Health under Award Number TL1 TR002318. **Author contributions:** A.B.R., C.M.R., R.A.M., and G.S. were mainly responsible for developing the method. A.B.R., C.M.R., R.A.M., A.K., P.S., S.M., W.C., and G.S. designed experiments. A.B.R., C.M.R., R.A.M., A.K., P.S., S.M., and W.C. performed experiments. Brain and spinal cord extraction was performed and supported by D.J.P., S.H.P., and D.L.S. A.B.R., C.M.R., Z.Y., and L.G. performed data analysis. Cluster annotation was performed by A.B.R., C.M.R., Z.Y., L.G., and B.T. A.B.R., C.M.R., B.T., and G.S. wrote the manuscript. **Competing interests:** A.R., R.M., and G.S. are inventors on a patent application (14/941,433) submitted by the University of Washington that covers the SPLiT-seq method. **Data availability:** All relevant sequencing files were deposited to GEO under ascension number GSE110823.

SUPPLEMENTARY MATERIALS

www.sciencemag.org/cgi/content/full/science.aam8999/DC1

Materials and Methods

Supplementary Text

Figs. S1 to S21

Tables S1 to S12

References (44 – 92)

1 February 2017; resubmitted 30 September 2017; accepted 26 February 2018

Published online 15 March 2018

10.1126/science.aam8999




Fig. 1. Overview of SPLiT-seq. (A) Labeling transcriptomes with split-pool barcoding. In each split-pool round, fixed cells or nuclei are randomly distributed into wells and transcripts are labeled with well-specific barcodes. Barcoded RT primers are used in the first round. Second and third round barcodes are appended to cDNA through ligation. A fourth barcode is added to cDNA molecules by PCR during sequencing library preparation. The bottom scheme shows the final barcoded cDNA molecule. (B) Species mixing experiment with a library prepared from 1,758 whole cells. Human UBCs are blue, mouse UBCs are red, and mixed-species UBCs are gray. The estimated barcode collision rate is 0.2%, whereas species purity is >99%. (C) UMI counts from mixing experiments performed with fresh and frozen (stored at -80°C for 2 weeks) cells and nuclei. Median human UMI counts for fresh cells: 15,365; frozen cells: 15,078; nuclei: 12,113; frozen nuclei: 13,636. (D) Measured gene expression by SPLiT-seq is highly correlated between frozen cells and cells processed immediately (Pearson- r : 0.987). Frozen and fresh cells were processed in two different SPLiT-seq experiments.




Fig. 2. Single-cell transcriptome landscape of postnatal brain and spinal cord development by SPLiT-seq. (A) Over 150,000 nuclei from P2 and P11 mouse brains and spinal cords were profiled in a single experiment employing over six million barcode combinations. Transcriptomes were clustered and then visualized using t-SNE. Cells are colored according to cell type. Each cluster was downsampled to 1,000 cells for visualization. (B) A total of 73 distinct clusters were assigned to nine cell classes based on expression of established markers. The violin plots show marker gene expression in each cluster. (C) Astrocyte clusters are highlighted in red in the t-SNE. The violin plots show markers that are differentially expressed between astrocyte subtypes. (D) Seven OPC and oligodendrocyte clusters (containing 10,087 nuclei) colocalized in the original t-SNE (highlighted in red), forming a lineage. Cells from these clusters were re-embedded with t-SNE. (E) The heatmap shows genes expressed differentially across pseudotime in the oligodendrocyte lineage.




Fig. 3. Neuronal clusters exhibit regional specificity. (A) Marker gene expression was used to map neuronal clusters to specific brain regions. (B) Sagittal composite RNA ISH maps for nine representative clusters from distinct areas. For each cell type, we averaged ISH intensities from the Allen DMBA across the top five differentially expressed genes. (C) Types of pyramidal neurons in the cortex display layer-specific enrichments according to marker genes: cortical pyramidal neurons are highlighted in red in the t-SNE. Expression of example marker genes in pyramidal clusters is shown in the middle and corresponding available RNA ISH results on the right. (D) Three clusters constitute a developmental trajectory in the hippocampus. Re-embedding these clusters highlights the branching of the two differentiation trajectories in pseudotime. (E) Expression of differentiation marker genes is overlaid on the t-SNE. RNA ISH maps (Allen DMBA) show the regional specificity of granule cell and pyramidal neuron markers.




Fig. 4. Neuronal differentiation trajectories in the cerebellum revealed by SPLiT-seq. (A) Major cell types and their locations in the cerebellum. (B) Two types of Purkinje cells with distinct gene expression programs were identified. Early Purkinje cells are primarily found in the P2 brain and late Purkinje cells in the P11 brain. (C) t-SNE re-embedding of 15,360 nuclei suggests a pseudotime ordering from proliferating, to migrating, to mature CGCs. (D) Expression of marker genes is overlaid on the t-SNE, and the corresponding RNA ISH from Allen DMBA is shown below. Marker genes associated with different layers of the cerebellum are expressed at different points in pseudotime. Gene expression order is consistent with ordering of the physical layers. RNA ISH maps confirm regional specificity of marker genes. (E) t-SNE re-embedding of 1,890 nuclei reveals a branching differentiation trajectory. Progenitors can either become Golgi cells or stellate/basket cells. (F) Markers for progenitors and mature cell types are expressed at different points in pseudotime and have layer specificity.




Fig. 5. Gene expression patterns and spatial origin of cell types in the spinal cord. (A) Re-clustering spinal cord nuclei resulted in 30 neuronal and 14 non-neuronal clusters. (B) GABAergic neurons were defined by expression of *Gad1* and *Gad2*. A subset of GABAergic neurons are also glycinergic, based on expression of *Slc6a5*. Glutamatergic neurons were defined by expression of VGLUT1 (*Slc17a6*), whereas cholinergic motor neurons express *Chat*. (C) Novel gene markers distinguish gamma motor neurons from alpha motor neurons. (D) Inferred spatial origin of neuronal clusters within the spinal cord. We analyzed the Allen Spinal Cord Atlas expression patterns of the top ten enriched genes in each cluster. Dark purple indicates expression of all ten genes in the given region, while white indicates none of the ten genes were expressed in the given region.

Single-cell profiling of the developing mouse brain and spinal cord with split-pool barcoding

Alexander B. Rosenberg, Charles M. Roco, Richard A. Muscat, Anna Kuchina, Paul Sample, Zizhen Yao, Lucas Gray, David J. Peeler, Sumit Mukherjee, Wei Chen, Suzie H. Pun, Drew L. Sellers, Bosiljka Tasic and Georg Seelig

published online March 15, 2018

ARTICLE TOOLS

<http://science.sciencemag.org/content/early/2018/03/14/science.aam8999>

SUPPLEMENTARY MATERIALS

<http://science.sciencemag.org/content/suppl/2018/03/14/science.aam8999.DC1>

REFERENCES

This article cites 88 articles, 23 of which you can access for free
<http://science.sciencemag.org/content/early/2018/03/14/science.aam8999#BIBL>

PERMISSIONS

<http://www.sciencemag.org/help/reprints-and-permissions>

Use of this article is subject to the [Terms of Service](#)